Continue



Directions: Answer these questions pertaining to the simplifying of radicals. Check your answer when finished. Assume any variables represent a positive quantity. 1. Expression: True or False a. b. c. d. 2. Expression: True or False a. b. c. d. 3. Expression: True or False a. b. c. d. Directions: Answer these questions pertaining to the simplifying of radicals. Check your answer when finished. Choose the best answer. The cartoon "hints" may, or may not, be helpful! 4. Simplify . 5. Which of the following cannot be simplified? 6. Which of the following is not equivalent to ? 7. Which of the following is true? 8. Simplify . NOTE: The re-posting of materials (in part or whole) from this site to the Internet is copyright violation and is not considered "fair use" for educators. Please read the "Terms of Use". Simplify the following radicals : Questions : 1) √202) √1213) √524) √455) 3√726) 3√407) 3√278) 4√2439) 5√28810) 6√320 Answers :1) 2√52) 113) 2√134) 3√55) 23√96) 23√57) 38) 3√39) 2√310) 2√5 Solution Practice questions on exponents and radicals Simplify the following : Practice questions on simplifying radicals with variables and exponents Simplify the following radicals with variables : Questions :1) √(16u4v3)2) √(147m3n3)3) √(75x2y)4) 6√(72x2)5) 3√(8x6y3)6) 3√(54m5n6)7) 4√(81m4n8)8) 5√(32p10q15) Answers :1) 4u2v√v2) 7mn√(3mn)3) 5x√(3y)4) 36x√25) 2x2y6) 3mn3√(2m2)7) 3mn28) 2p2q3 Solution Practice questions on operations with radicals Simplify the following radical expression : 1) 7√30 + 2√75 + 5√50 Solution2) √27 + √105 + √108 + √45 Solution3) √45 + 3√20 + √80 - 4√40 Solution4) 3√5 + 2√95 + 3√117 - √78 Solution5) 3 √32 - 2√8 + √50 Solution6) 2 √12 - 3√27 - √243 Solution7) √54 - √2500 - √24 Solution8) √45 - √25 - √80 Solution9) 5√95 - 2√50 - 3√180 Solution10) √5 x √18 Solution11) ∛7 x ∛8 Solution12) ∛13 x ∛5 Solution13) 3√35 ÷ 2√7 Solution14) 15√54 ÷ 3√6 Solution15) (48)1/4 ÷ (72)1/8 Solution16) 4√3, 18√2, -3√3, 15√2 Solution17) 2∛2, 24∛2, - 4∛2 Solution Answers :1) √30 + 5 √3 + 5 √2 2) 4 √3 + √1053) 9 √5 - 2 √104) 3 √5 + 2 √95 + 9 √13 - √785) 13 √26) -14 √37) -√6 -508) -√5 - 59) 5 √95 - 10 √2 - 18 √510) 3 √511) 2 ∛712) ∛6513) (3/2)√514) 1515) (32)1/816) √3 + 33√217) 22 ∛2 Practice questions on rationalizing the denominator with surds Rationalize the denominator : Practice questions on rationalizing the denominator with variables Rationalize the denominator :1) 1/√x Solution2) 1/(x + √y) Solution3) (√x + y) / (x - √y) Solution4) (√x + √y)/√x Solution5) (√x + √y)/(√x - √y) Solution6) 1/(x + y√3) Solution7) 3√(2/3a) Solution8) Simplify : √(12x2) / √(30x) Solution9) √(100x/11y) Solution Answers :1) √x/x2) (x - √y) / (x2 - y)3) [x√x + √xy + xy + y√y] / (x2 - y2)4) [x + √(xy)]/x5) (x + 2√(xy) + y) / (x - y)6) (x - y√3) / (x2 + 3y2)7) 3√(18a2) / 3a8) √10x / 59) 10√(11xy) / 11y Practice questions on using distributive property with radicals Expand and simplify the following radicals : Questions :1) √2 (√5 + √2)2) √3 (1 - √3)3) √11 (2√11 - 1)4) 2√3 (√3 - √5)5) (1 + √2) (2 + √2)6) (√3 + 2) (√3 - 1)7) (√5 + 2) (√5 - 3)8) (2√2 + √3) (2√2 - √3)9) (2 + √3) (2 + √3)10) (4 - √2) (3 + √2)11) (√7 - √3) (√7 + √3)12) (4 - √2) (3 - √2) Answers :1) √10 + 22) √3 - 3 3) 22 - √114) 6 - 2√155) 4 + 3√26) 1 + √37) -1 - √58) 59) 7 + 4√310) 10 + √2 11) 412) 14 - 7√2 Solution Practice questions on solving for unknown with radicals Solve for unknown variables :1) (x + y√2) (2 - √2) = 1+√22) (2 - 3√2)(x + y√2) = √23) (x + y√2) (3 + √2) = 14) (a + √2) (2 - √2) = 4 - b√25) (a + b√2)2 = 33 + 20√26) (x + y√2) (3 - √2) = -4√2 Answers :1) x = 2 and y = 3/22) x = -3/7 and y = -1/74) a = 3 and b = - 15) a = 5 and b = 26) x = -8/7 and y = -12/7 Solution Problem 1 :2√3(4 - √3)SolutionProblem 2 :- √2(2 - √3)SolutionProblem 3 :2√3(√3 - 1) - 2√3SolutionProblem 4 :(2√2 - 5)(1 - √2)SolutionProblem 5 :(3 + 2√5)(2 - √5)SolutionProblem 6 :(4 - √2)(3 + 2√2)SolutionProblem 7 :(3 - √7)2SolutionProblem 8 :- (2 - √5)2SolutionProblem 9 :(2 - √3)(2 + √3)SolutionProblem 10 :(5 - √3)(5 + √3)SolutionProblem 11 :Write √1/7 in the form k√7SolutionProblem 12 :Find x, y ∈ Q such that (3 + x√5) (√5 – y) = - 13 + 5√5SolutionProblem 13 :Find p, q ∈ Q such that (p + 3√7) (5 + q√7) = 9√7 - 53SolutionProblem 14 :Solve for m, √(m - 1) + 5 = m - 2Solution 1) 8√3 – 62) 2 - 2√23) 6 - 4√34) 7√2 - 95) √5 - 46) 8 + 5√27) 16 - 6√78) 4√5 - 99) 110) 2211) 1/712) x = - 2 and y = 113) q = 10/21 and q = -314) m = 10 and m = 5 Kindly mail your feedback to v4formath@gmail.comWe always appreciate your feedback. ©All rights reserved. onlinemath4all.com Home » Arithmetic » Simplifying Radicals Worksheets Show Mobile Notice Show All Notes Hide All Notes Mobile Notice You appear to be on a device with a "narrow" screen width (i.e. you are probably on a mobile phone). Due to the nature of the mathematics on this site it is best viewed in landscape mode. If your device is not in landscape mode many of the equations will run off the side of your device (you should be able to scroll/swipe to see them) and some of the menu items will be cut off due to the narrow screen width. For problems 1 – 4 write the expression in exponential form. For problems 5 – 7 evaluate the radical. For problems 8 – 12 simplify each of the following. Assume that x, y and z are all positive. \(\sqrt[3]{{{x^8}}}\) Solution \(\sqrt {8{y^3}} \) Solution \(\sqrt[4]{{{x^7}{y^{20}}{z^{11}}}}\) Solution \(\sqrt[3]{{54{x^6}{y^7}{z^2}}}\) Solution \(\sqrt[4]{{4{x^3}y}}\,\,\sqrt[4]{{8{x^2}{y^3}{z^5}}}\) Solution For problems 13 – 15 multiply each of the following. Assume that x is positive. \(\sqrt x \left({4 - 3\sqrt x } \right)\) Solution \(\left({2\sqrt x + 1} \right)\left({3 - 4\sqrt x } \right)\) Solution \(\left({\sqrt[3]{x} + 2\,\,\sqrt[3]{{{x^2}}}} \right)\left({4 - \sqrt[3]{{{x^2}}}} \right)\) Solution For problems 16 – 19 rationalize the denominator. Assume that x and y are both positive. \(\displaystyle \frac{6}{{\sqrt x }}\) Solution \(\displaystyle \frac{9}{{\sqrt[3]{{2x}}}}\) Solution \(\displaystyle \frac{4}{{\sqrt x + 2\sqrt y }}\) Solution \(\displaystyle \frac{{10}}{{3 - 5\sqrt x }}\) Solution ← Previous 1 2 3 4 5 Next → Which of the following answer choices best simplifies ? Possible Answers: Correct answer: Explanation: The first step to simplifying a problem like this one is to convert all radicals to fractional exponents. Remember the following relationship: Also keep in mind your exponent rules, especially this one: Now, let's get started on this problem. First, we change that radical expression into something with fractional exponents instead. Now we use our exponent rules to simplify the numerator. Finally, we simplify the entire fraction: We can leave it like this, but it would be better to write it this way, without negative exponents: Simplify: You may assume that the radicand is nonnegative. Possible Answers: Correct answer: Explanation: The polynomial , being a polynomial of degree 2, cannot be a cube of another polynomial. Also, it does not fit the pattern of a perfect square binomial, since its constant term is negative. Therefore, we cannot extract a root of the polynomial to help to simplify it. We can, however, rewrite each root as a fractional exponent, apply the power of a power property, then convert back, as follows: Simplify: You may assume that is a nonnegative real number. Possible Answers: Correct answer: Explanation: The best way to simplify a radical within a radical is to rewrite each root as a fractional exponent, then convert back. First, rewrite the roots as exponents. We can simplify this further: What is the value of ? Possible Answers: Correct answer: Explanation: Recall that when you have a fractional exponent, this means that you have a root involved. The denominator of the exponent is the type of root. Our question's exponent is: Therefore, the root is or a square root. The numerator is the power for the base. Therefore, we can rewrite our problem as: Now, to simplify this, we could do: Using our exponent rules, this is: Factoring out sets of , we get: , or What is the value of ? Possible Answers: Correct answer: Explanation: Recall that when you have a fractional exponent, this means that you have a root involved. The denominator of the exponent is the type of root. Our question's exponent is: Therefore, the root is or a cube root. The numerator is the power for the base. Therefore, we can rewrite our problem as: Now, to simplify this, we could do: Using our exponent rules, this is: Factoring out sets of and set of , we get: What is the value of ? Possible Answers: Correct answer: Explanation: Recall that when you have a fractional exponent, this means that you have a root involved. The denominator of the exponent is the type of root. Our question's exponent is: Therefore, the root is or a cube root. The numerator is the power for the base. Therefore, we can rewrite our problem as: Now, to simplify this, we could do: We can factor out a set of . This leaves us with: Simplifying, this is: Which of the following is equivalent to ? Possible Answers: Correct answer: Explanation: Recall that the cube root of a number is the number that when multiplied by itself 3 times, yields your number. Thus, we want y, where: Consider That cannot be our solution. Then try Thus, this must be our solution. Next time, remember that radicals can be represented by fractional exponents! Choose the best answer. Reduce the following in exponential form: Possible Answers: Correct answer: Explanation: Simplify the inner term within the parentheses. Simplify, Possible Answers: Correct answer: Explanation: First write the square root of as an exponent, From the rules of exponents we know we can simplyfy by adding the exponents, Simplify: Possible Answers: Correct answer: Explanation: To simplify the expression, we must remember that a fraction as a power denotes a radical: the numerator is the power to which the term is taken inside the radical, and the denominator denotes the degree of the root (i.e. 2 means square root, 3 means cube root, etc.) Rewriting our expression, we get which expanded becomes Now, move the cubes outside of the cube root, after taking their cube root, leaving behind the terms that aren't cubes: ← Previous 1 2 3 4 5 Next → Robert Certified Tutor Abilene Christian University, Bachelor of Science, Mathematics. Lipscomb University, Masters in Education, Education. Isaias Certified Tutor Instituto Tecnológico de Merida (ITM), Bachelor of Science, Biochemical Engineering. Yucatan Center for Scientific Research (... Nafis Certified Tutor University of Leipzig, Germany., Bachelor of Science, Physics. University of Pittsburgh-Pittsburgh Campus, Doctor of Philosop... If you've found an issue with this question, please let us know. With the help of the community we can continue to improve our educational resources. If you believe that content available by means of the Website (as defined in our Terms of Service) infringes one or more of your copyrights, please notify us by providing a written notice ("Infringement Notice") containing the information described below to the designated agent listed below. If Varsity Tutors takes action in response to an Infringement Notice, it will make a good faith attempt to contact the party that made such content available by means of the most recent email address, if any, provided by such party to Varsity Tutors. Your Infringement Notice may be forwarded to the party that made the content available or to third parties such as ChillingEffects.org. Please be advised that you will be liable for damages (including costs and attorneys' fees) if you materially misrepresent that a product or activity is infringing your copyrights. Thus, if you are not sure content located on or linked-to by the Website infringes your copyright, you should consider first contacting an attorney. Please follow these steps to file a notice: You must include the following: A physical or electronic signature of the copyright owner or a person authorized to act on their behalf; An identification of the copyright claimed to have been infringed; A description of the nature and exact location of the content that you claim to infringe your copyright, in \ sufficient detail to permit Varsity Tutors to find and positively identify that content; for example we require a link to the specific question (not just the name of the question) that contains the content and a description of which specific portion of the question – an image, a link, the text, etc – your complaint refers to; Your name, address, telephone number and email address; and A statement by you: (a) that you believe in good faith that the use of the content that you claim to infringe your copyright is not authorized by law, or by the copyright owner or such owner's agent; (b) that all of the information contained in your Infringement Notice is accurate, and (c) under penalty of perjury, that you are either the copyright owner or a person authorized to act on their behalf. Send your complaint to our designated agent at: Charles Cohn Varsity Tutors LLC 101 S. Hanley Rd, Suite 300 St. Louis, MO 63105 Or fill out the form below: Mathworksheetsgo.com is now a part of Mathwarehouse.com. All of your worksheets are now here on Mathwarehouse.com. Please update your bookmarks! This printable PDF worksheet is designed for students in grades 5-8 to master simplifying radical expressions. The worksheet features 25 scaffolded questions, progressing from easy to challenging, along with model problems and a comprehensive answer key for effective learning and self-assessment. Simplifying radicals is a fundamental concept in algebra, essential for students in grades 5 through 8. It involves expressing a radical in its simplest form by removing any perfect square factors from under the radical sign. This process not only makes the radical easier to understand and work with, but it also lays the groundwork for more advanced algebraic manipulations. The core idea behind simplifying radicals is to identify hidden perfect squares within the radicand (the number under the radical) and extract their roots. For instance, √75 can be simplified because 75 contains a perfect square factor, 25 (25 x 3 = 75). By recognizing this, we can rewrite √75 as √(25 x 3) and then simplify it to 5√3. Mastering this skill is crucial as it appears in various areas of mathematics, including solving equations, working with geometric figures, and performing calculus operations. Our comprehensive worksheet provides numerous practice problems to solidify understanding and build confidence in simplifying radicals. With a step-by-step approach, students will learn to break down complex radicals into their simplest forms, unlocking the beauty and simplicity of algebraic expressions. Understanding Perfect Squares Before diving into the process of simplifying radicals, it's crucial to have a solid grasp of perfect squares. A perfect square is an integer that can be obtained by squaring another integer. In simpler terms, it's the result of multiplying a whole number by itself. Examples of perfect squares include 1 (1×1), 4 (2×2), 9 (3×3), 16 (4×4), 25 (5×5), and so on. Recognizing these numbers is the first step towards simplifying radicals effectively. Understanding perfect squares allows us to identify factors within the radicand that can be simplified. When we encounter a radical like √36, knowing that 36 is a perfect square immediately tells us that √36 simplifies to 6. However, not all radicands are perfect squares themselves. In such cases, we need to find perfect square factors within the radicand. For example, when simplifying √20, we recognize that 20 can be factored into 4 x 5, where 4 is a perfect square. Therefore, we can rewrite √20 as √(4 x 5) and then simplify it to 2√5. This ability to identify and extract perfect square factors is what makes simplifying radicals possible. Our worksheet provides ample practice in recognizing and working with perfect squares, laying a strong foundation for mastering radical simplification. Factoring to Find Perfect Squares The key to simplifying radicals often lies in the ability to factor the radicand (the number under the radical sign) and identify perfect square factors. This process involves breaking down the radicand into its constituent factors, searching for factors that are perfect squares. For instance, when simplifying √48, we need to find the largest perfect square that divides evenly into 48. One approach is to start by listing the factors of 48: 1, 2, 3, 4, 6, 8, 12, 16, 24, and 48. Among these factors, we identify 16 as the largest perfect square (4×4). Therefore, we can rewrite √48 as √(16 x 3). This allows us to simplify the radical as follows: √48 = √(16 x 3) = √16 x √3 = 4√3. Alternatively, you can use prime factorization to break down the radicand into its prime factors. For example, the prime factorization of 48 is 2 x 2 x 2 x 2 x 3, which can be grouped as (2 x 2) x (2 x 2) x 3 = 4 x 4 x 3 = 16 x 3. This again leads us to recognize 16 as the perfect square factor. Practice with various examples in our worksheet will help you develop fluency in factoring and identifying perfect square factors, making radical simplification a straightforward process. Simplifying Radical Expressions: Step-by-Step Simplifying radical expressions becomes manageable when approached systematically. The first step involves identifying whether the radicand (the expression under the radical) has any perfect square factors. If it does, you can proceed to simplify; if not, the radical is already in its simplest form. Next, factor the radicand to isolate the perfect square. For example, to simplify √75, recognize that 75 can be factored into 25 x 3, where 25 is a perfect square (5×5). Thus, √75 becomes √(25 x 3). Now, apply the product property of radicals, which states that √(ab) = √a x √b. Using this property, we can rewrite √(25 x 3) as √25 x √3. Then, simplify the perfect square root: √25 = 5. The expression now becomes 5√3, which is the simplified form of √75. For more complex expressions, repeat these steps until the radicand contains no more perfect square factors. Always double-check your work to ensure that you have extracted the largest possible perfect square. Practicing consistently with our worksheets will solidify your understanding of these steps. Adding and Subtracting Radical Expressions Adding and subtracting radical expressions requires a foundational understanding of simplifying radicals. The key principle is that you can only combine "like radicals," meaning radicals with the same radicand (the number under the radical symbol) and the same index (the root being taken, such as square root or cube root). Before attempting to add or subtract, simplify each radical expression individually. This involves factoring out perfect squares (or perfect cubes, etc.) as demonstrated in previous sections. Once each radical is simplified, examine the terms to identify like radicals. For instance, consider the expression 3√2 + 5√2. Here, both terms have the same radicand (2) and the same index (square root). Therefore, they are like radicals and can be combined. To do so, simply add or subtract their coefficients (the numbers in front of the radical): 3 + 5 = 8. The result is 8√2. If the radicals are not alike, you cannot directly combine them. For example, 2√3 + 4√5 cannot be simplified further because √3 and √5 are different radicals. However, sometimes simplification reveals like radicals that were not initially apparent. Combining Like Terms in Radical Expressions Combining like terms is a fundamental algebraic skill that extends seamlessly to radical expressions. The core concept remains the same: only terms with identical variable parts can be combined. In the context of radicals, "like terms" refers to radical expressions that have the same radicand (the expression under the radical symbol) and the same index (the root being taken). Before combining any terms, it's crucial to simplify each radical expression individually. This often involves factoring out perfect squares (or cubes, etc.) and applying the rules of radical simplification discussed earlier. Simplifying first can reveal hidden like terms that were not initially obvious. Once the radical expressions are simplified, identify the like terms. These are the terms that have the exact same radical portion. For example, in the expression 5√x + 3√x 2√y, the terms 5√x and 3√x are like terms because they both contain √x. The term -2√y is not a like term because it contains √y, which is different from √x. To combine like terms, simply add or subtract their coefficients (the numerical part in front of the radical). In our example, 5√x + 3√x = (5+3)√x = 8√x. Therefore, the simplified expression would be 8√x 2√y. Simplifying Radicals with Variables Simplifying radicals containing variables builds upon the principles of simplifying numerical radicals, incorporating the rules of exponents. When a variable is raised to a power under a radical, the goal is to extract any perfect square (or cube, etc.) factors. For example, in √(x²), x² is a perfect square, and its square root is x. If the exponent of the variable under the radical is even, the simplification is straightforward. Divide the exponent by the index of the radical. The result is the exponent of the variable outside the radical. For instance, √(x⁴) = x^(4/2) = x². If the exponent is odd, separate the variable into two factors: one with the highest possible even exponent and another with an exponent of 1. Consider √(x⁵). We can rewrite this as √(x⁴ * x). Then, √(x⁴) simplifies to x², leaving us with x²√x. The variable x with the even exponent (x⁴) is extracted from the radical, while the remaining variable x stays inside. When simplifying radicals with multiple variables, apply the same process to each variable independently. For example, √(x³y⁴) becomes √(x² * x * y⁴) = xy²√x. The absolute value is sometimes necessary when extracting variables from radicals, particularly when the index is even and the variable's original exponent is odd, to ensure the result is non-negative. Rationalizing Denominators Rationalizing the denominator is a process used to eliminate radicals from the denominator of a fraction. This is typically done to simplify expressions and make them easier to work with. The basic idea is to multiply both the numerator and denominator of the fraction by a suitable expression that will eliminate the radical in the denominator. When the denominator contains a single term with a square root, you multiply both the numerator and the denominator by that square root. For example, to rationalize 1/√2, you multiply both the numerator and denominator by √2, resulting in √2 / 2. If the denominator is a binomial containing a square root, such as (a + √b) or (a √b), you multiply both the numerator and denominator by the conjugate of the denominator. The conjugate is formed by changing the sign between the two terms. For example, the conjugate of (2 + √3) is (2 √3). Multiplying by the conjugate utilizes the difference of squares pattern: (a + b)(a b) = a² b². This eliminates the square root from the denominator. For instance, to rationalize 1/(2 + √3), you multiply both the numerator and denominator by (2 √3), which results in (2 √3) / (4 3) = 2 √3. Free Worksheet (PDF) Availability A free, printable worksheet in PDF format is readily available to assist students in mastering the simplification of radicals. This valuable resource offers a comprehensive set of practice problems designed to reinforce the concepts and techniques involved in simplifying radical expressions. The worksheet is easily accessible and can be downloaded for convenient use at home or in the classroom. The PDF format ensures that the worksheet maintains its formatting and clarity across various devices and operating systems. This makes it a versatile tool for students using computers, tablets, or smartphones. The worksheet is designed to be printer-friendly, allowing for easy duplication and distribution. Teachers and parents can utilize this free worksheet as a supplementary resource to enhance their lessons or provide additional practice opportunities. The availability of the worksheet promotes accessibility to quality educational materials, supporting students in their learning journey. The worksheet covers a range of difficulty levels, making it suitable for students with varying levels of understanding. The included answer key further enhances the worksheet's value by enabling students to check their work and identify areas where they may need additional support. This self-assessment feature promotes independent learning and helps students develop a deeper understanding of simplifying radicals. Answer Key and Solutions An answer key is provided with the simplifying radicals worksheet, offering complete solutions to each problem. This feature allows students to check their work and gain immediate feedback on their understanding of the concepts. The answer key serves as a valuable tool for self-assessment and independent learning. By comparing their solutions to the provided answers, students can identify any errors or misconceptions they may have. The detailed solutions in the answer key not only provide the correct answers but also demonstrate the step-by-step process of simplifying each radical expression. This allows students to understand the reasoning behind each step and develop a deeper understanding of the underlying mathematical principles. The answer key is designed to be user-friendly and easy to navigate, making it simple for students to find the solutions they need. It is organized in a clear and concise manner, ensuring that students can quickly locate the answers to specific problems. Teachers and parents can also benefit from the answer key, as it provides a convenient way to assess student progress and identify areas where additional instruction may be needed. The answer key can be used as a guide for providing targeted support and ensuring that students are mastering the essential skills of simplifying radicals. It promotes effective learning and helps students build confidence in their mathematical abilities. Scaffolded Questions: Easy to Challenging The simplifying radicals worksheet features a carefully designed set of 25 scaffolded questions, progressing in difficulty from easy to challenging. This structure allows students to gradually build their understanding and skills in simplifying radical expressions. The initial questions focus on basic concepts, such as identifying perfect squares and simplifying simple radicals. As students progress through the worksheet, the questions become increasingly complex, requiring them to apply their knowledge to more challenging problems. These later questions may involve simplifying radicals with variables, rationalizing denominators, or combining like terms in radical expressions. The scaffolded approach ensures that students are not overwhelmed by the complexity of the material. By starting with easier questions, students can build confidence and develop a solid foundation in the fundamental concepts. As they progress to more challenging questions, they are able to apply their knowledge and skills in a more sophisticated way. This carefully structured progression makes the worksheet suitable for students of varying skill levels. Students who are new to simplifying radicals can start with the easier questions and gradually work their way up to the more challenging ones. Students who have already mastered the basic concepts can focus on the more challenging questions to further develop their skills. Applications of Simplifying Radicals Simplifying radicals is not just an abstract mathematical exercise; it has numerous practical applications in various fields. In geometry, simplifying radicals is essential when calculating the lengths of sides in right triangles using the Pythagorean theorem, especially when dealing with irrational numbers. For instance, finding the diagonal of a square with side length 'a' involves simplifying √2a² to a√2. In physics, simplifying radicals arises in calculations involving energy, momentum, and wave functions. Many physical formulas contain square roots, and simplifying these expressions can lead to more manageable and interpretable results. Engineering disciplines also utilize simplified radicals when dealing with structural analysis, signal processing, and control systems. Furthermore, simplifying radicals is crucial in computer graphics and game development. Calculating distances, angles, and transformations often involves square roots, and simplifying these expressions can improve computational efficiency and reduce rendering times. Even in fields like finance, simplifying radicals can be useful when analyzing growth rates and compound interest. Understanding how to simplify radicals allows students to solve real-world problems and appreciate the practical relevance of this mathematical concept beyond the classroom. For each problem, simplify, and then check your answers and record your total score below. Example: 4√49 By the end of this section, you will be able to: Use the Product Property to simplify radical expressions Use the Quotient Property to simplify radical expressions Before you get started, take this readiness quiz. We will simplify radical expressions in a way similar to how we simplified fractions. A fraction is simplified if there are no common factors in the numerator and denominator. To simplify a fraction, we look for any common factors in the numerator and denominator. A radical expression, \(\sqrt[n]{a}\), is considered simplified if it has no factors of \(m^{n}\). So, to simplify a radical expression, we look for any factors in the radicand that are powers of the index. For real numbers \(a\) and \(m\), and \(n\geq 2\), \(\sqrt[n]{a}\) is considered simplified if \(a\) has no factors of \(m^{n}\) For example, \(\sqrt{5}\) is considered simplified because there are no perfect square factors in \(5\). But \(\sqrt{12}\) is not simplified because \(12\) has a perfect square factor of \(4\). Similarly, \(\sqrt[3]{4}\) is simplified because there are no perfect cube factors in \(4\). But \(\sqrt[3]{24}\) is not simplified because \(24\) has a perfect cube factor of \(8\). To simplify radical expressions, we will also use some properties of roots. The properties we will use to simplify radical expressions are similar to the properties of exponents. We know that \[(a b)^{n}=a^{n} b^{n}.\] The corresponding of Product Property of Roots says that \[\sqrt[n]{a b}=\sqrt[n]{a} \cdot \sqrt[n]{b}.\] If \(\sqrt[n]{a}\) and \(\sqrt[n]{b}\) are real numbers, and \(n\geq 2\) is an integer, then \(\sqrt[n]{a b}=\sqrt[n]{a} \cdot \sqrt[n]{b} \quad \text { and } \quad \sqrt[n]{a} \cdot \sqrt[n]{b}=\sqrt[n]{a b}\) We use the Product Property of Roots to remove all perfect square factors from a square root. Simplify: \(\sqrt{98}\). Solution: Step 1: Find the largest factor in the radicand that is a perfect power of the index. We see that \(49\) is the largest factor of \(98\) that has a power of \(2\). \(\sqrt{98}\) Rewrite the radicand as a product of two factors, using that factor. In other words \(49\) is the largest perfect square factor of \(98\). \(98 = 49\cdot 2\) Always write the perfect square factor first. \(\sqrt{49\cdot 2}\) Step 2: Use the product rule to rewrite the radical as the product of two radicals. \(\sqrt{49} \cdot \sqrt{2}\) Step 3: Simplify the root of the perfect power. \(7\sqrt{2}\) Simplify: \(\sqrt{48}\) Answer \(4 \sqrt{3}\) Simplify: \(\sqrt{45}\). Answer \(3 \sqrt{5}\) Notice in the previous example that the simplified form of \(\sqrt{98}\) is \(7\sqrt{2}\), which is the product of an integer and a square root. We always write the integer in front of the square root. Be careful to write your integer so that it is not confused with the index. The expression \(7\sqrt{2}\) is very different from \(\sqrt[7]{2}\). Find the largest factor in the radicand that is a perfect power of the index. Rewrite the radicand as a product of two factors, using that factor. Use the product rule to rewrite the radical as the product of two radicals. Simplify the root of the perfect power. We will apply this method in the next example. It may be helpful to have a table of perfect squares, cubes, and fourth powers. Simplify: \(\sqrt{500}\) \(\sqrt[3]{16}\) \(\sqrt[4]{243}\) Solution: a. \(\sqrt{500}\) Rewrite the radicand as a product using the largest perfect square factor. \(\sqrt{100 \cdot 5}\) Rewrite the radical as the product of two radicals. \(\sqrt{100} \cdot \sqrt{5}\) Simplify. \(10\sqrt{5}\) b. \(\sqrt[3]{16}\) Rewrite the radicand as a product using the greatest perfect cube factor. \(2^{3}=8\) \(\sqrt[3]{8 \cdot 2}\) Rewrite the radical as the product of two radicals. \(\sqrt[3]{8} \cdot \sqrt[3]{2}\) Simplify. \(2 \sqrt[3]{2}\) c. \(\sqrt[4]{243}\) Rewrite the radicand as a product using the greatest perfect fourth power factor. \(3^{4}=81\) \(\sqrt[4]{81 \cdot 3}\) Rewrite the radical as the product of two radicals. \(\sqrt[4]{81} \cdot \sqrt[4]{3}\) Simplify. \(3 \sqrt[4]{3}\) Simplify: a. \(\sqrt{288}\) b. \(\sqrt[3]{81}\) c. \(\sqrt[4]{64}\) Answer a. \(12\sqrt{2}\) b. \(3 \sqrt[3]{3}\) c. \(2 \sqrt[4]{4}\) Simplify: a. \(\sqrt{432}\) b. \(\sqrt[3]{625}\) c. \(\sqrt[4]{729}\) Answer a. \(12\sqrt{3}\) b. \(5 \sqrt[3]{5}\) c. \(3 \sqrt[4]{9}\) The next example is much like the previous examples, but with variables. Don't forget to use the absolute value signs when taking an even root of an expression with a variable in the radical. Simplify: \(\sqrt{x^{3}}\) \(\sqrt[3]{x^{4}}\) \(\sqrt[4]{x^{7}}\) Solution: a. \(\sqrt{x^{3}}\) Rewrite the radicand as a product using the largest perfect square factor. \(\sqrt{x^{2} \cdot x}\) Rewrite the radical as the product of two radicals. \(\sqrt{x^{2}} \cdot \sqrt{x}\) Simplify. \(|x| \sqrt{x}\) b. \(\sqrt[3]{x^{4}}\) Rewrite the radicand as a product using the largest perfect cube factor. \(\sqrt[3]{x^{3} \cdot x}\) Rewrite the radical as the product of two radicals. \(\sqrt[3]{x^{3}} \cdot \sqrt[3]{x}\) Simplify. \(x \sqrt[3]{x}\) c. \(\sqrt[4]{x^{7}}\) Rewrite the radicand as a product using the greatest perfect fourth power factor. \(\sqrt[4]{x^{4} \cdot x^{3}}\) Rewrite the radical as the product of two radicals. \(\sqrt[4]{x^{4}} \cdot \sqrt[4]{x^{3}}\) Simplify. \(|x| \sqrt[4]{x^{3}}\) Simplify: a. \(\sqrt{b^{5}}\) b. \(\sqrt[4]{y^{6}}\) c. \(\sqrt[3]{z^{5}}\) Answer a. \(b^{2} \sqrt{b}\) b. \(|y| \sqrt[4]{y^{2}}\) c. \(z \sqrt[3]{z^{2}}\) Simplify: a. \(\sqrt{p^{9}}\) b. \(\sqrt[5]{y^{8}}\) c. \(\sqrt[6]{q^{13}}\) Answer a. \(p^{4} \sqrt{p}\) b. \(p \sqrt[5]{p^{3}}\) c. \(q^{2} \sqrt[6]{q}\) We follow the same procedure when there is a coefficient in the radicand. In the next example, both the constant and the variable have perfect square factors. Simplify: \(\sqrt{72 n^{7}}\) \(\sqrt[3]{24 x^{7}}\) \(\sqrt[4]{80 y^{14}}\) Solution: a. \(\sqrt{72 n^{7}}\) Rewrite the radicand as a product using the largest perfect square factor. \(\sqrt{36 n^{6} \cdot 2 n}\) Rewrite the radical as the product of two radicals. \(\sqrt{36 n^{6}} \cdot \sqrt{2 n}\) Simplify. (6\left|n^{3}\right| \sqrt{2 n}\) b. \(\sqrt[3]{24 x^{7}}\) Rewrite the radicand as a product using perfect cube factors. \(\sqrt[3]{8 x^{6} \cdot 3 x}\) Rewrite the radical as the product of two radicals. \(\sqrt[3]{8 x^{6}} \cdot \sqrt[3]{3 x}\) Rewrite the first radicand as \((2 x^{2}\right)^{3}\). \(\sqrt[3]{\left(2 x^{2}\right)^{3}} \cdot \sqrt[3]{3 x}\) Simplify. (2 x^{2} \sqrt[3]{3 x}\) c. \(\sqrt[4]{80 y^{14}}\) Rewrite the radicand as a product using perfect fourth power factors. \(\sqrt[4]{16 y^{12} \cdot 5 y^{2}}\) Rewrite the radical as the product of two radicals. \(\sqrt[4]{16 y^{12}} \cdot \sqrt[4]{5 y^{2}}\) Rewrite the first radicand as \((2 y^{3}\right)^{4}\). \(\sqrt[4]{\left(2 y^{3}\right)^{4}} \cdot \sqrt[4]{5 y^{2}}\) Simplify: a. \(\sqrt{32 y^{5}}\) b. \(\sqrt[3]{54 p^{10}}\) c. \(\sqrt[4]{64 q^{10}}\) Answer a. \(4 y^{2} \sqrt{2 y}\) b. \(3 p^{3} \sqrt[3]{2 p}\) c. \(2 q^{2} \sqrt[4]{4 q^{2}}\) Simplify: a. \(\sqrt{75 a^{9}}\) b. \(\sqrt[3]{128 m^{11}}\) c. \(\sqrt[4]{162 n^{7}}\) Answer a. \(5 a^{4} \sqrt{3 a}\) b. \(4 m^{3} \sqrt[3]{2 m^{2}}\) c. \(3|n| \sqrt[4]{2 n^{3}}\) In the next example, we continue to use the same methods even though there are more than one number under the radical. Simplify: \(\sqrt{63 u^{3} v^{5}}\) \(\sqrt[3]{40 x^{4} y^{5}}\) \(\sqrt[4]{48 x^{4} y^{7}}\) Solution: a. \(\sqrt{63 u^{3} v^{5}}\) Rewrite the radicand as a product using the largest perfect square factor. \(\sqrt{9 u^{2} v^{4} \cdot 7 u v}\) Rewrite the radical as the product of two radicals. \(\sqrt{9 u^{2} v^{4}} \cdot \sqrt{7 u v}\) Rewrite the first radicand as \(\left(3 u v^{2}\right)^{2}\). \(\sqrt{\left(3 u v^{2}\right)^{2}} \cdot \sqrt{7 u v}\) Simplify. \(3|u| v^{2} \sqrt{7 u v}\) b. \(\sqrt[3]{40 x^{4} y^{5}}\) Rewrite the radicand as a product using the largest perfect cube factor. \(\sqrt[3]{8 x^{3} y^{3} \cdot 5 x y^{2}}\) Rewrite the radical as the product of two radicals. \(\sqrt[3]{8 x^{3} y^{3}} \cdot \sqrt[3]{5 x y^{2}}\) Rewrite the first radicand as \((2xy)^{3}\). \(\sqrt[3]{(2 x y)^{3}} \cdot \sqrt[3]{5 x y^{2}}\) Simplify. (2 x y \sqrt[3]{5 x y^{2}}\) c. \(\sqrt[4]{48 x^{4} y^{7}}\) Rewrite the radicand as a product using the largest perfect fourth power factor. \(\sqrt[4]{16 x^{4} y^{4} \cdot 3 y^{3}}\) Rewrite the radical as the product of two radicals. \(\sqrt[4]{16 x^{4} y^{4}} \cdot \sqrt[4]{3 y^{3}}\) Rewrite the first radicand as \((2xy)^{4}\). \(\sqrt[4]{(2 x y)^{4}} \cdot \sqrt[4]{3 y^{3}}\) Simplify. \(2|x y| \sqrt[4]{3 y^{3}}\) Simplify: \(\sqrt{98 a^{7} b^{5}}\) \(\sqrt[3]{56 x^{5} y^{4}}\) \(\sqrt[4]{32 x^{5} y^{8}}\) Answer \(7\left|a^{3}\right| b^{2} \sqrt{2 a b}\) \(2 x y \sqrt[3]{7 x^{2} y}\) \(2|x| y^{2} \sqrt[4]{2 x}\) Simplify: \(\sqrt{180 m^{9} n^{11}}\) \(\sqrt[3]{72 x^{6} y^{5}}\) \(\sqrt[4]{80 x^{7} y^{4}}\) Answer \(6 m^{4}\left|n^{5}\right| \sqrt{5 m n}\) \(2 x^{2} y \sqrt[3]{9 y^{2}}\) \(2|x| y \sqrt[4]{5 x^{3}}\) Simplify: \(\sqrt[3]{-27}\) \(\sqrt[4]{-16}\) Solution: a. \(\sqrt[3]{-27}\) Rewrite the radicand as a product using perfect cube factors. \(\sqrt[3]{(-3)^{3}}\) Take the cube root. \(-3\) b. \(\sqrt[4]{-16}\) There is no real number \(n\) where \(n^{4}=-16\). Not a real number Simplify: \(\sqrt[3]{-64}\) \(\sqrt[4]{-81}\) Answer Simplify: \(\sqrt[3]{-625}\) \(\sqrt[4]{-324}\) Answer \(-5 \sqrt[3]{5}\) no real number We have seen how to use the order of operations to simplify some expressions with radicals. In the next example,